GARY MANGIACOPA ARCHIVE
==========
Title: B2F1I6
Slug: b2f1i6
Categories: Uncategorized
Source: https://garymangiacopraarchive.com/b2f1i6
Pages: 10 scanned, 10 extracted
OCR: Google Vision API (document_text_detection)
Processed: 2026-06-06
==========

THE FAR SIDE. 1982. Universal Press Syndicate, Reprinted with permission. All rights reserved.

NATURE

The larger-than-life lizard

Geckos are familiar to most people as the sticky-footed lizards that lurk behind the lights in buildings in warmer climes. Apart from their distinctive shape and ability to walk up walls and windows, one feature that the many species in the family Gekkonidae have in common is their small size. Thus, in July 1984, an Iranian medical orderly made herpetological history when he came across a gecko measuring 40cm in length. Mohamed Reza Ensaf made his discovery in Khuzistan, the troubled border region near Iraq, and was subsequently immortalised by having the species, Eublepharis ensafi, named after him. It was believed to be the largest gecko on earth (beating the former contender, the 30cm tokay, by a third)-but it was about to be dwarfed by a zoological mystery.

The ink was scarcely dry on the above description when a century-old, stuffed lizard moved into the limelight.

Alain

Delcourt, curator of the Marseille Natural History Museum in France, sent some photographs of this puzzling specimen to George Zug, herpetologist at the Smithsonian
12
Institution,
p.12
BBC Wildlife, Bristol, E(1:1), January 1988, p.s
Giant gecko Hoplodactylus delcourti, resurrected for the record books.
Washington DC, for possible identification. He in turn forwarded the data to Aaron Bauer, of the University of Berkeley, California, and Anthony Russell, of the University of Calgary, Canada, who described the animal as a new species of gecko, and named it Hoplodactylus delcourti in honour of the Marseilles museum curator.
For a gecko, this species is truly monstrous-62cm (more than two feet) in length-twice as long as a tokay, and three times the length of the former
"You Imbecilel... We flew 12,000 miles for THIS?"
largest species of Hoplodactylus. Its colour is a yellowish-brown with dark, reddish-brown longitudinal dorsal stripes-evidently a spectacular creature when alive-but no details of the specimen's origin are known today. It was probably brought to the Marseille Museum sometime during the years 1833 to 1869-a period for which the archives have been lost-and it is astonishing to think that visitors to the museum have been shown the stuffed enigma for a century and a half without realising its zoological significance!
The genus Hoplodactylus is endemic to New Zealand; is it possible that a giant stripy gecko

could have eluded antipodean naturalists for more than two centuries? And did the Maoris know of the species? Although the largest known reptile in New Zealand is the tuatara, which reaches only 60cm, Maori tribesmen told James Cook in 1777 of eight foot (two and a half metre) lizards that "sometimes seize and devour men".

Similar tales of huge saurians called taniwha have been recorded by other explorers, and it has been suggested that they may refer to either saltwater crocodiles or monitor lizards- neither of which have been recorded by science in New Zealand. But these are clearly not referring to geckos.

There are, however, other reports which do fit the bill. In the Transactions of the New Zealand Institute of 1871, Sir Walter Buller wrote about the lizards of New Zealand, including a mysterious one: "The Kawekaweau, a beautiful striped lizard, sometimes attaining a length of two feet, is still undescribed. It was formerly abundant in the forests north of Auckland, and is still occasionally met with. Mr F E Maning, of Hokianga, recently obtained possession of a pair of live ones, but unfortunately for science, one of them was devoured by a cat and the other made its escape.

In 1873, a Major W G Mair wrote in the same journal of the similar sounding kawekau, which a Maori chief had killed under loose bark. The chief described it as being "about two foot long, and as thick as a man's wrist; colour brown, striped longitudinally with dull red." These descriptions both match closely the Marseille museum specimen, and the

reported behaviour (hidden under bark) is consistent with the known daytime retreats of other large geckos. Bauer and Russell have thus proposed that Hoplodactylus delcourti and the kawekau (alias kawekaweau) of the Maoris are one and the same species.

Another mysterious reptile which may be a gecko, but which doesn't fit with the above reports, is the ngarara, of which there are descriptions of at least two varieties: one is larger and lived in burrows on the plains, the other was smaller and lived in streams. A report in the Auckland Weekly News of 11 April 1956 described a creature that may have been an aquatic ngarara. And it does not seem beyond the bounds of possibility that a nocturnal forest gecko-even a giant stripy one-might have eluded the attentions of collectors in some remote corner of New Zealand. Anyone visiting that part of the world who notices such a creature stomping out from behind the wall light should contact their nearest natural history immediately!

museum

MICHEL RAYNAL

BBC WILDLIFE JANUARY 1988

m d'Histoire Naturelle de Marseille, France

[PAGE BREAK]

J. Ethnobiol. 7(1):83-91

To Michel Raynal,
wishes
with best
Summer 1987

HOPLODACTYLUS DELCOURTI (REPTILIA:GEKKONIDAE) AND THE KAWEKAWEAU OF MAORI FOLKLORE

AARON M. BAUER
Department of Zoology and Museum of Vertebrate Zoology
University of California
Berkeley, CA 94720

and
ANTHONY P. RUSSELL
Department of Biology
The University of Calgary
2500 University Drive N.W.
Calgary, Alberta, Canada T2N 1N4
and
Department of Ichthyology and Herpetology
Royal Ontario Museum
100 Queen's Park
Toronto, Ontario, Canada M5S 2C6
ABSTRACT.-The recent discovery of a giant gecko of unknown provenance in the collections of the Musée d'Histoire Naturelle de Marseille necessitated the employment of relatively unorthodox sources in order to attempt to establish its identity. A combination of morphological characteristics and anecdotal information from folkloric sources assisted in establishing the potential geographic locality of the new species. This determination indicated that an animal heretofore thought to be purely mythical, the kawekaweau of Maori folklore, is indeed a recognizable biological species.

INTRODUCTION

A hitherto unknown species of giant gecko was recently described (Bauer and Russell 1986) from a specimen rediscovered among the collections of the Musée d'Histoire Naturelle de Marseille, France. It has been in the collections for at least 117 years (Bauer and Russell 1986). The specimen has been assigned to the genus Hoplodactylus, a taxon of live-bearing geckos distributed throughout New Zealand but known from nowhere else. Strong anatomical evidence supports this placement of the new species-Hoplodactylus delcourti. Although no locality data are available for the specimen, we believe that anecdotal evidence, much of it dating from the last century, may be used to associate this giant gecko with the kawekaweau, a large reptile of Maori folklore (Bauer and Russell 1986). This therefore links the specimen with New Zealand. The evidence we present serves as an example of the value of native folkloric sources in certain aspects of zoological alpha systematics.

THE NEW ZEALAND GEKKONID FAUNA

The native New Zealand gekkonid fauna consists of two major lineages-the nocturnal "brown geckos" (genus Hoplodactylus) and the diurnal "green geckos" (genus Naultinus). (We consider Heteropholis Fischer 1883 to be subsumed within Naultinus-see Thomas [1982]). Both genera are members of the tribe Carphodactylini within the subfamily Diplodactylinae, a group restricted in its distribution to Australia, New Zealand

[PAGE BREAK]

and New Caledonia (Fig. 1). Within this group, the New Zealand geckos have their closest affinities with the New Caledonian genera Rhacodactylus, Bavayia and Eurydactylodes (Bauer 1986). In particular, all share a similar arrangement of preanal pores on the venter and a strongly clawed first digit with scansorial pad. Although no uniquely derived characters support the monophyly of Hoplodactylus (Bauer 1986), digital characters and scalation allow members of this genus to be distinguished from other carphodactyline geckos.

HOPLODACTYLUS DELCOURTI, ITS AFFINITIES AND PROBABLE PROVENANCE

Hoplodactylus delcourti is distinguished from other members of the genus chiefly

by its huge size (370 mm head and body length [SVL]). This makes the giant Hoplodactylus by far the largest known gecko species (Bauer and Russell 1986). It is 54% larger than the next largest species (Rhacodactylus leachianus) and more than 130% larger than the largest known individual of Hoplodactylus duvaucelii (Whitaker 1968), the next largest congener. The significance of and possible origins of such great size are discussed by Russell and Bauer (1986). Whether its point of origin was New Zealand or New Caledonia, Hoplodactylus delcourti represents a trend toward island gigantism that has been widely documented in reptiles (Mertens 1934; Carlquist 1974) as well as other vertebrates and invertebrates. New Caledonia, for example, is home to the second and fourth largest extant geckos in the world as well as to one of the largest skinks, a giant grasshopper and a huge flying fox. New Zealand, of course, is noted for the tallest bird, Dinornis giganteus, (Cracraft 1976), numerous large skinks and a group of large orthopterans (wetas). The causes of island gigantism are unclear.

FIG. 1.-Distribution of the tribe Carphodactylini is restricted to Australia, New Zealand and New Caledonia (left map). The kawekaweau has been reported from Waimana (Mair 1873) and possibly from the Waoku Plateau (Walsh 1905) and Wanganui (Downes 1937), all on the North Island of New Zealand (right map).

[PAGE BREAK]

FIG. 2.-An example of Maori wood carving depicting a lizard-like animal (perhaps a gecko) with
paua (abalone) eyes. This specimen is in the collections of the National
Museum of New Zealand in Wellington.

Large lizards known to the Maori were the mokonui (Kerry-Nichols 1884; Best 1923), the kumi (Hector 1899) (believed by Hardy [1977] to be mythical), the ngarara (a 450 mm lizard from Kaiapoi in North Canterbury, Stack 1875) and the kaweau of kawekaweau reported from several areas of the North Island. All available reports of these lizards seem to be secondhand, and the characters attributed to the animals are

often incongruous. In particular, the reference to the aquatic habits of the kawekaweau is surely inconsistent with all geckos and most of the known skinks of New Zealand. Indeed, many of the accounts of early settlers may refer to Australian agamid lizards (Hector 1899), such as Physignathus lesueuri, a large water-frequenting lizard.

ANECDOTAL SOURCES IN RELATION TO A NEW ZEALAND ORIGIN

A number of the descriptions of the lizards of Maori legend offer corroborating evidence as to the identity and origin of Hoplodactylus delcourti. The most valuable information comes from Mair (1873) who commented on "the existence of a large forest lizard, called by the Maoris kaweau". (The spelling and pronunciation of this term varies among Maori tribal dialects-Best 1909.) He continues, "In 1870 an Urewera chief

[PAGE BREAK]

Morphological features clearly place the animal in the genus Hoplodactylus, but this does not provide information about its origin. The morphology of the specimen limits the areas of possible origin, as it represents a member of a limited radiation of the tribe Carphodactylini in the southwest Pacific Ocean. One possibility is that the animal originated in the French island territory of New Caledonia. A New Caledonian origin would more easily explain the presence of the specimen in a French museum but would require that the genus Hoplodactylus, in the recent past, occurred in areas other than New Zealand. An alternative hypothesis, that of a New Zealand origin for Hoplodactylus delcourti, would be consistent with current ideas regarding the distribution of this genus, and would not be inconsistent with the history of many areas of the North Island. French explorers, scientists and settlers were quite active in the North Island, particularly in the area of the Bay of Islands (Hocken 1908; Wright 1950). While the museum in Marseille does not possess any other herpetological specimens from New Zealand, it does possess a number of old specimens from Australia and other areas in the Pacific. The dating of the acquisition of the specimen is problematic, but it is most likely that it was acquired between 1833 and 1869. No accessionary documentation of the museum survives from this period. The only New Caledonian material currently in the collec-[AD] tion dates from the period 1902-1905 and represents donations from the Marseille Zoological Park, which had received a number of New Caledonian lizards and sea snakes from M. Bernier of the Muséum de Nouméa (Vayssière 1916).

REPTILES AND MAORI FOLKLORE

The contention that the specimen is from New Zealand is also supported by historical and anecdotal records from both Maori and pakeha (European) sources. Maori knowledge of natural history in New Zealand has been gained in their 900 year occupancy of these islands, but is tempered by common Polynesian conceptions and terminology of nature brought with them from their prior experiences in the islands of the central Pacific (McLintock 1966). There is considerable anecdotal material suggesting the existence of undescribed vertebrates in New Zealand. Of particular note is the case of the "New Zealand otter" (Watson 1960) which has never been resolved. Allusion to unknown reptiles of various sizes and attributes have also been made frequently in the local folklore and lizards, or lizard-like animals feature prominently in Maori art (Best 1923; Skinner 1964. See Fig. 2). Cook (1777) was the first European to record the Maori tales of giant reptiles and many authors have reiterated Maori claims of such beasts (Jameson 1842; Buller 1878, Gisborne 1888; Best 1909). Some of these animals, such as the taniwha, a giant reptile-like creature, have been summarily dismissed because of the great sizes credited to them. Yet, even in these claims there may be some veracity by way of the correlation of reports of large lizards with crocodiles or monitors (family Varanidae)

encountered by the ancestors of the Maoris or collateral groups of Polynesians in other areas of the Pacific (Downes 1937). Lizard-like reptiles of a smaller size also figure prominently in Maori legend. Many of these tales can almost certainly be referred to the tuatara (Sphenodon), but others seem to describe one or more different animals.
The folk taxonomy of the Maori, like that of many peoples, seems to be quite exact (Best 1909), with most of the currently recognized taxa being identified by unique Maori names. Numerous early European workers provided partial listings of Maori names for local animals (e.g. Buller 1878, Downes 1937). Among the herpetofauna of New Zealand, it seems that only the leiopelmatid frogs and some of the more remotely distributed geckos of the South Island were completely unknown to the Maoris. Lumping of biological species also occurred among the skinks, but this is understandable given that modern herpetologists can only distinguish some of the morphologically similar Leiolopisma species on the basis of biochemical features (C. Daugherty, pers. comm.).
55
85

[PAGE BREAK]

with nocturnal habits, isolated and hidden retreats, and low population density might go unnoticed in New Zealand today.
Whether now extinct or not, Hoplodactylus delcourti has not occurred in large numbers for at least a century. Two human factors may be suggested to have contributed to this decline. The first is the degradation of primary habitat through the destruction of native forests. A second is the introduction and spread of introduced mammals by both the Maori and later pakeha settlers of New Zealand. Stack (1875) suggested rats were responsible for the disappearance of a 450 mm ngarara from the Waimakariri River Valley, while Buller (1895) cited feral pigs as being the most destructive predators of the kawekaweau. Taylor (1868) and Kirk (1895) add cats to the list of introduced enemies of reptiles.
Rats have long been regarded as being severely damaging to populations of native reptiles in New Zealand. The effect of rats on the tuatara has been extensively documented (Crook 1973a, 1973b, 1975; Newman 1983). Although it is currently unclear whether the main factor involved is predation, competition or degradation of habitat, it is known that islands supporting the kiore (Rattus exulans) have fewer juvenile tuataras and
adult recruitment than do similar islands without rats. Whitaker (1973, 1978;
poorer
Bull and Whitaker 1975) documented that the presence of rats decreases both lizard population size and species diversity on offshore islands. His assumption was that the primary danger posed by rats is predation, especially on young lizards. This view is supported by empirical evidence showing that rats do ingest lizards (Bettesworth 1972; Bettesworth and Anderson 1972; Whitaker 1978). Whitaker (1978) drew conclusions about vulnerability which are pertinent to Hoplodactylus delcourti: a) nocturnal species are more susceptible to rats than are diumal species, and b) larger species are more vulnerable than are smaller ones. In the former case activity time corresponds with peak rat activity, and in the latter animals are unable to use hiding places inaccessible to rats. These conclusions clearly place the kawekaweau at very high risk.
While no certainty can be attached to the origin of the giant gecko, a body of evidence from historical and ethnographic sources suggests that the species did or does inhabit

parts of the North Island. The likelihood of the evolution of such a gigantic form in New Zealand is supported by the preponderance of large species of a variety of taxonomic groups (ratites, scincids, orthopterans, etc.).

ANECDOTAL SOURCES AND THEIR UTILITY IN ALPHA SYSTEMATICS

Common local animals, as well as rarer but more spectacular ones, are strongly associated with the art, folklore and traditions of most cultures. As shown by Maori folklore and legend, as recorded by European chroinclers, the knowledge of native peoples about the fauna may be of value to systematic zoologists. In this particular case the attributes of a relatively rare animal were still known to Maoris who had never seen the animal alive. This is probably due in part to the large size of the kawekaweau, but also to the fact that mythical powers had been associated with the animal and it therefore took on more symbolic significance. Best (1909:236; 1923:331) stated that it represented the souls of dead forbears of any person who saw it and indicated it was time for that individual to join these ancestors in the spirit world.

Some generalities may be outlined for the use of such anecdotal native evidence in the examination of systematic questions. To the zoologist many native references seem implausible because they attribute biologically incorrect (or impossible) characteristics to animals. In the case of the taniwha (Downes 1937), virtually all of the attributes of the beast are entirely mythical. The combination of plausible features with mythical ones may, however, imply that the animal in question is based in fact.

[PAGE BREAK]

killed one under the loose bark of a dead rata, in the Waimana Valley [Fig. 1], he described it to me as being about two feet long and as thick as a man's wrist; color brown, striped longitudinally with dull red". This description, one of the few to provide morphological information about giant lizards, matches extremely well the size and color of the specimen at Marseille, and the position of the lizard under bark is not inconsistent with the known daytime retreats of other large geckos. Buller (1895) stated that the kawekaweau was semi-arboreal and existed in the deep forests of the North Island at least until the 1860s. He also commented on the striking "banded" pattern of the animal. Best (1909, 1923) referred to the kawekaweau as being the size of a small tuatara, reddish in color, and to its retreats as being the holes of trees. This again is a likely gekkonid hiding place and is supported by the known retreats of giant geckos of the related genus Rhacodactylus (Meier 1979).

Kerry-Nichols (1884) claimed that a large lizard (ngarara in the ethnotaxonomic generic sense) was known to local Maoris to inhabit the caves and rock piles of the North Island. Walsh (1905) remarked on an unnamed large lizard, with both arboreal and aquatic habits from the Waoku Plateau (Fig. 1), between the Hokianga and Kaipara districts, the remains of which washed down the Waima Creek around 1870. Finally, Jameson (1842) referred to a large lizard resembling the chameleon. It is noteworthy that in New Caledonia, home of the previously largest known geckos (Rhacodactylus), the local name for large geckos is "cameleons", distinguishing them from the small house geckos (Hemidactylus and Lepidodactylus) which are known as "margouillats". It seems that large geckos are frequently confused with chameleons in areas where chameleons do not occur. This is probably because some species in each family change color, or perhaps because of the prominence of the eyes and relatively slow movements of the animals.

Hardy (1977) suggested that the skink Leiolopisma gracilicorpus, might be identifiable as the kawekaweau. However, the only known specimen of this species has a length

of 97 mm and is thus far smaller than the legendary kawekaweau. Further, the specimen is completely bleached, so that a correlation with the color pattern described in the early accounts is not possible. Given the existence of a specimen matching features of color and morphology as reported by early New Zealanders, we feel that the new species of giant gecko may indeed be the basis for the legendary kawekaweau.

POTENTIAL AGENTS OF EXTINCTION IN HISTORICAL TIMES

Although the lack of material and records would indicate that Hoplodactylus delcourti has become extinct, this is by no means certain. There remain in the North Island extensive areas of relatively undisturbed land which might harbor remaining populations. These include areas containing kauri forests as well as rock outcrops, two of the possible habitats of the kawekaweau. It is certainly no less likely that this animal still survives than it is that the much larger thylacine (Thylacinus cynocephalus) of Tasmania is extant. A number of vertebrate species long thought extinct have been "rediscovered" in the recent past. These include the Chaco peccary (Catagonus c.f. wagneri) (Wetzel et al. 1975), Leadbeater's possum (Gymnobelideus leadbeateri) and New Zealand's own takahe (Notornis mantelli). The takahe (witness the Maori name) as well as at least some species of Moa were present as distinct entities in Maori folk taxonomy. Although most, if not all, the species of Moa had become extinct by the time Europeans had begun to anthropologically document Maori culture, knowledge of the existence of these giant birds had been maintained by oral tradition in at least some tribes (Andrews 1986). Even more noteworthy are recently discovered vertebrates about which nothing was previously known. Two of the smaller, but nonetheless spectacular congeners of H. delcourti, H. rakiurae Thomas 1981 and H. kahutarae Whitaker 1984 have only recently been discovered in remote parts of New Zealand. It is just possible that a 370 mm SVL gecko

[PAGE BREAK]

Features such as size are particularly likely to be exaggerated. Consequently, reports of animals of relatively small size or rather unspectacular appearance are also more likely to be referrable to living creatures. Unfortunately, rare animals such as the kawekaweau, had seldom been seen by those describing them. Thus these animals are generally reported as occuring "several villages away" and to have been seen by "a relative" or "acquaintance". Thus again, the fact that some of the Maoris responsible for the claims reported to the pakeha, whose reports are cited herein, actually saw the animals themselves, near their own villages and in the recent past, adds credibility to the factual basis of the animal in question.

The scientific identification of animals largely from folkloric sources is rare, but such an instance is applicable in the case of Hoplodactylus delcourti. In such instances there must always remain a degree of uncertainty of identification. Nonetheless, the reciprocal illumination of the fields of zoology and ethnology holds great potential for uncovering useful information. Greenwell (1985) has reviewed the categorization of cryptozoological evidence and has provided an operational classificatory system for documenting cases. The current case falls somewhere between Greenwell's (1985) categories III and IV. Here we have a taxon represented by a partial specimen and "known" only from anecdotal sources. It is precisely these sources that have aided in its identification.

ACKNOWLEDGMENTS

The authors thank Garth Underwood and Alain Delcourt for bringing the existence of this specimen to their attention. A. H. Whitaker, B. W. Thomas, D. Towns, B. Gill, G. Hardy, D. G. Newman and J. Robb assisted in locating obscure references and in following up leads on the specimen
and its associated folklore. G. M. Christman prepared the map. The insightful comments of two referees (A. Rea and W. Sherbrooke) assisted greatly in helping us to clarify certain concepts. Ms. Jane Larson typed the manuscript.

LITERATURE CITED

ANDREWS, J.R.H. 1986. The Southern Ark. Zoological Discovery in New Zealand [AD] 1769-1900. Century Hutchinson, Auckland.

BAUER, A. M. 1986. Systematics, biogeography and evolutionary morphology of the Carphodactylini (Reptilia:Gekkonidae). Unpubl. Ph.D. Thesis (Zoology) Univ. of California, Berkeley.

and A. P. RUSSELL. 1986. Hoplodactylus delcourti n. sp. (Reptilia Gekkonidae), the largest known gecko. New Zealand J. Zool. 13:141-148.

BEST, E. 1909. Maori forest lore: being some account of native forest lore and woodcraft and also of many myths, rites, customs and superstitions connected with the flora and fauna of the Tuhoe or Ure-wera district. Part 2. Trans. New Zealand Inst. 41:231-285.

1923. Notes on the occurrence of the lizard in Maori carvings. New Zealand J. Sci. Tech. 5(6):321-335.

BETTESWORTH, D. J. 1972. Rattus exulans on Red Mercury Island. Tane 18:117-118.

and G. R. V. ANDERSON. 1972. Diet of Rattus norvegicus on Whale Island, Bay of Plenty, New Zealand. Tane 18: 189-195.

BULL, P. C. and A. H. WHITAKER. 1975. The amphibians, reptiles, birds and mammals. Pp. 231-276 in Biogeography and Ecology in New Zealand. Kuschel, G. (ed.), Junk, The Hague.

BULLER, J. 1878. Forty years in New Zealand: Including a personal narrative, an account of Maoridom, and of the Christianization and colonization of the country. Hodder and Stoughton, London.

BULLER, W. 1895. Illustrations of Darwinism; or, the avifauna of New Zealand considered in relation to the fundamental Law of Descent with modification. Trans. New Zealand Inst. 27:75-104.

CARLQUIST, S. 1974. Island Biology. Columbia University Press, New York.

[PAGE BREAK]

LITERATURE CITED (continued)
COOK, J. A. 1777. A voyage toward the South Pole and round the world. 2 Volumes. London.
CRACRAFT, J. 1976. The species of Moas (Aves:Dinomithidae). Smithsonian Contrib. to Paleobiol. 27:189-205.
CROOK, I. G. 1973a. The tuatara, Sphenodon punctatus Gray, on islands with and without populations of the Polynesian rat, Rattus exulans (Peale). Proc. New Zealand Ecol. Soc. 20:115-120.
1973b. Tuatara and the Polynesian rat. Wildlife-a review 4:47-48.
1975. The tuatara. Pp. 331-352 in Biogeography and Ecology in New Zealand. Kuschel, G. (ed.), Junk, The Hague.
DOWNES, T. W. 1937. Maori mentality regarding the lizard and taniwha in the Whanganui River area. J. Polynesian Soc. 46(184):206-224.
GISBORNE, W. 1888. The Colony of New Zealand: Its history, vicissitudes and progress. Petherich & Co., London.
GREENWELL, J. R. 1985. A classificatory scheme for cryptozoology. Cryptozoology. 4:1-14.
HARDY, G. S. 1977. The New Zealand Scincidae (Reptilia:Lacertilia), a taxonomic and zoogeographic study. New Zealand J. Zool. 4:221-325.
HECTOR, J. 1899. On the kumi. Trans. New Zealand Inst. 31:717-718,720.
HOCKEN, DR. 1908. Early visits of the French to New Zealand. Trans. New Zealand Inst. 40:137-153.
JAMESON, R. G. 1842. New Zealand, South Australia and New South Wales: a record of recent travels in these colonies, with special reference to emigration and the advantages employment of labour and capital. Smith, Elder, and Co., London.
KERRY-NICHOLS, J. H 1884. The King

Country; or, explorations in New Zealand. A narrative of 600 miles of travel through Maoriland. Sampson, Low, Marston, Searle, & Rivington, London.
KIRK, T. 1895. The displacement of species in New Zealand. Trans. New Zealand Inst. 28:1-27.
MAIR, W. G. 1873. Notes on Rurima Rocks. Trans. New Zealand Inst. 5:151-153.
MCLINTOCK, A. H. (ed.). 1966. An Encyclopedia of New Zealand. 3 Vols. R. E. Owen, Wellington.
MEIER, H. 1979. Herpetologische Beobachtungen auf Neukaledonien. Salamandra 15:113-139.
MERTENS, R. 1934. Die Insel-Reptilien, ihre Ausbreitung, Variation und Artbildung. Zoologica (Stuttgart) 32(84):1-209.
MILLENER, P. R. 1981. The Quaternary avifauna of the North Island, New Zealand. Two volumes. Unpubl. Ph.D. Thesis (Zoology) Univ. Auckland.
NEWMAN, D. G. (ed.). 1982. New Zealand Herpetology Proceedings of a Symposium held at Victoria University of Wellington, January 1980. New Zealand Wildlife Service Occ. Pub. No. 2, Wellington.
1983. Tuatara/kiore relationships. Wildlife-a review 12:60-63.
RUSSELL, A. P. and A. M. BAUER. 1986. Le gecko geant Hoplodactylus delcourti et ses relations avec le gigantisme et l'endemisme insulaire chez les Gekkonidae. Mesogee 46:25-28.
SKINNER, H. D. 1964. Crocodile and lizard in New Zealand. Myth and material culture. Rec. Otago Mus. Anthropology No. 1.
STACK, J. W. 1875. On the disappearance of the larger kinds of lizards from North Canterbury. Trans. New Zealand Inst. 7:295-297.
TAYLOR, R. 1868(?). Essay on the zoology of New Zealand. MS (Alexander Tumbull Library, Wellington).
THOMAS, B. W. 1982. New Zealand Gekkonidae: Evolutionary trends and taxa. Pp. 333-339 in Newman, D. G. (ed.) New Zealand Herpetology. New Zealand Wildlife Service Occ. Pub. No. 2.
VAYASSIÈRE, A. 1916. Rapport sur la Section de Zoologie du Muséum. Ann. Mus.

d'Hist. Nat. Marseille 15:1-7.
WALSH, P. 1905. A rare saurian. Trans. New Zealand Inst. 37:351-352.
WATSON, J. S. 1960. The New Zealand "otter". Rec. Canterbury Mus. 7(3):175-183.
WETZEL, R. M., R. E. DUBOS, R. L. MARTIN and P. MYERS. 1975. Catagonus, an "extinct" peccary, alive in Paraguay. Science 189:379-381.
WHITAKER, A. H. 1968. The lizards of the Poor Knights Islands, New Zealand. New Zealand J. Sci. 11(4):623-651.
1973. Lizard populations on islands with and without Polynesian rats,

[PAGE BREAK]

UG
THE
UNIVERSITY
OF CALGARY
2500 University Drive N.W., Calgary, Alberta, Canada T2N 1N4
FACULTY OF SCIENCE
Department of BIOLOGICAL SCIENCES
[AD] Telephone (403) 220-5261
July 29, 1988
M. Michel Raynal
Tour Rubis
Apartement C24
36 Avenue d'Italie
75013 Paris
FRANCE
Dear M. Raynal,
Thank you very much for your letter of May 02, 1988. I am sorry for the delay in replying. I have been busy with local field work this summer. On August 04 I am leaving for New Zealand and Australia for three weeks. The main reason for going to New Zealand is to attempt to search out Hoplodactylus delcourti. We have some good leads and local contacts, and appreciate people such as yourself keeping us informed of reports or possible reports of this animal. We will be submitting the results of our search to "Cryptozoology", and hope, of course, that they include some tangible evidence of the real thing.
I will be in contact again on my return. Thank you for your continued interest and encouragement.
Sincerely,
Wirel
Anthony P. Russell
APR:sps
Olympic Village and Speedskating - 1988